

## COVID-19 Results Briefing

### Vermont

December 15, 2022

This document contains summary information on the latest projections from the IHME model on COVID-19 in Vermont. The model was run on December 15, 2022, with data through December 7, 2022.

### Current situation

- Daily infections in the last week decreased to 10,000 per day on average compared to 11,000 the week before (Figure 1.1).
- Daily reported cases in the last week increased to 72 per day on average compared to 49 the week before (Figure 2.1).
- Daily hospital census in the last week (through December 7) increased to 52 per day on average compared to 34 the week before.
- Reported deaths due to COVID-19 in the last week increased to two per day on average compared to one the week before (Figure 3.1).
- Total deaths due to COVID-19 in the last week increased to seven per day on average compared to four the week before (Figure 3.1). This makes COVID-19 the number one cause of death in Vermont this week (Table 1). Estimated total daily deaths due to COVID-19 in the past week were 4.2 times larger than the reported number of deaths.
- The daily rate of reported deaths due to COVID-19 is greater than 4 per million in one state (Figure 4.1).
- The daily rate of total deaths due to COVID-19 is greater than 4 per million in two states (Figure 4.2).
- We estimate that 98% of people in Vermont have been infected at least once as of December 12 (Figure 6.1). Effective R, computed using cases, hospitalizations, and deaths, is greater than 1 in 41 states and the District of Columbia. Effective R in Vermont was 1.6 on December 1 (Figure 7.1).
- The infection-detection rate in Vermont was close to 3% on December 12 (Figure 8.1).
- Based on the GISAID and various national databases, combined with our variant spread model, we estimate the current prevalence of variants of concern (Figures 9.1-9.6). We estimate that the Alpha variant is circulating in 22 states and the District of Columbia, that the Beta variant is circulating in no states nor the District of Columbia, that the Delta variant is circulating in 50 states and the District of Columbia, that the Gamma variant is circulating in no states nor the District of Columbia, that the BA.1/BA.2 variants are circulating in 50 states and the District of Columbia, and that the BA.5 variant is circulating in 50 states and the District of Columbia.

### Trends in drivers of transmission

- Based on self-reported mask use data collected in the COVID-19 Trends and Impact Survey, an estimated 4% of people are projected to always wear a mask when leaving their home. Mask use after June 24, 2022 is a statistical forecast.
- As of December 12, 36 states and the District of Columbia have reached 70% or more of the population who have received at least one vaccine dose, and 24 states and the District of Columbia have reached 70% or more of the population who are fully vaccinated (Figures 12.1 and 12.2). 94% of people in Vermont have received at least one vaccine dose, and 88% are fully vaccinated.
- In our current reference scenario, we expect that 591,000 people will be vaccinated with at least one dose by April 1 (Figure 14.1). We expect that 89% of the population will be fully vaccinated by April 1.

### Projections and scenarios

We produce three scenarios when projecting COVID-19. The **reference scenario** is our forecast of what we think is most likely to happen:

- Vaccines are distributed at the expected pace. Brand- and variant-specific vaccine efficacy is updated using the latest available information from peer-reviewed publications and other reports.
- Future mask use will decline to 50% of the minimum level it reached between January 1, 2021, and May 1, 2022. This decline begins after the last observed data point in each location and transitions linearly to the minimum over a period of six weeks.
- Mobility increases as vaccine coverage increases.
- Mandates will be reimposed at the maximum level of mandates in the post-ancestral period once the death rate has reached an algorithmic minimum threshold of daily reported deaths for a given location.
- 80% of those who are fully vaccinated (two doses for most vaccines, or one dose for Johnson & Johnson) receive an additional dose six months after becoming fully vaccinated, and 80% of those who receive an additional dose receive a second additional dose six months later.
- Antiviral utilization for COVID-19 risk prevention has reached 80% in high-risk populations and 50% in low-risk populations between March 1, 2022, and June 1, 2022. This applies in high-income countries, but not low- and middle-income countries, and this rollout assumption follows a similar pattern to global vaccine rollouts.

The **80% mask use scenario** makes all the same assumptions as the reference scenario but assumes all locations reach 80% mask use within seven days. If a location currently has higher than 80% use, mask use remains at the current level.

The **antiviral access scenario** makes all the same assumptions as the reference scenario but assumes globally distributed antivirals and extends coverage to all low- and middle-income countries between August 15, 2022, and September 15, 2022.

### Infections

- Daily estimated infections in the **reference scenario** will decline to 370 by February 16, 2023 (Figure 16.1).
- Daily estimated infections in the **80% mask use scenario** will decline to 150 by February 24, 2023 (Figure 16.1).
- Daily estimated infections in the **antiviral access scenario** will decline to 370 by February 16, 2023 (Figure 16.1).

### Cases

- Daily estimated cases in the **reference scenario** will rise to 370 by December 21, 2022 (Figure 16.2).
- Daily estimated cases in the **80% mask use scenario** will rise to 350 by December 20, 2022 (Figure 16.2).
- Daily estimated cases in the **antiviral access scenario** will rise to 370 by December 21, 2022 (Figure 16.2).

### Hospitalizations

- Daily hospital census in the **reference scenario** will rise to 250 by December 28, 2022 (Figure 16.3). At some point from December through April 1, four states will have high or extreme stress on hospital beds (Figure 18.1). At some point from December through April 1, one state will have high or extreme stress on intensive care unit (ICU) capacity (Figure 19.1).
- Daily hospital census in the **80% mask use scenario** will rise to 220 by December 27, 2022 (Figure 16.3).
- Daily hospital census in the **antiviral access scenario** will rise to 250 by December 28, 2022 (Figure 16.3).

### Deaths

- In our **reference scenario**, our model projects 930 cumulative reported deaths due to COVID-19 on April 1. This represents 140 additional deaths from December 12 to April 1. Daily reported COVID-19



deaths in the **reference scenario** will rise to zero by December 28, 2022 (Figure 16.4).

- Under our **reference scenario**, our model projects 3,900 cumulative total deaths due to COVID-19 on April 1. This represents 580 additional deaths from December 12 to April 1 (Figure 16.5).
- In our **80% mask use scenario**, our model projects 920 cumulative reported deaths due to COVID-19 on April 1. This represents 120 additional deaths from December 12 to April 1. Daily reported COVID-19 deaths in the **80% mask use scenario** will rise to zero by December 28, 2022 (Figure 16.4).
- In our **antiviral access scenario**, our model projects 930 cumulative reported deaths due to COVID-19 on April 1. This represents 140 additional deaths from December 12 to April 1. Daily reported COVID-19 deaths in the **antiviral access scenario** will rise to zero by December 28, 2022 (Figure 16.4).
- Figure 17.1 compares our reference scenario forecasts to other publicly archived models. Forecasts are widely divergent.

## Model updates

We have updated our reference scenario to assume that mandates will be re-imposed at the maximum level of mandates in the post-ancestral period once the death rate has reached an algorithmic minimum threshold of daily reported deaths for a given location.


For the foreseeable future, we will not be updating our model or producing COVID-19 estimates. These will be the final briefing documents we produce until further notice.

**Figure 1.1: Daily COVID-19 hospital census and estimated infections****Figure 2.1: Reported daily COVID-19 cases, moving average**

**Table 1: Ranking of total deaths due to COVID-19 among the leading causes of mortality this week, assuming uniform deaths of non-COVID causes throughout the year**

Cause name	Weekly deaths	Ranking
COVID-19	49	1
Ischemic heart disease	23	2
Tracheal, bronchus, and lung cancer	9	3
Chronic obstructive pulmonary disease	9	4
Stroke	7	5
Alzheimer’s disease and other dementias	6	6
Colon and rectum cancer	4	7
Chronic kidney disease	3	8
Diabetes mellitus	3	9
Falls	3	10

**Figure 3.1: Smoothed trend estimate of daily COVID-19 deaths**



Daily COVID-19 death rate per 1 million on December 12, 2022

Figure 4.1: Daily reported COVID-19 death rate per 1 million





Figure 4.2: Daily total COVID-19 death rate per 1 million



Cumulative COVID-19 deaths per 100,000 on December 12, 2022

Figure 5.1: Reported cumulative COVID-19 deaths per 100,000






Figure 5.2: Total cumulative COVID-19 deaths per 100,000







**Figure 6.1: Estimated percent of the population infected with COVID-19 on December 12, 2022**



**Figure 7.1: Mean effective R on December 1, 2022.** Effective R less than 1 means that transmission should decline, all other things being held the same. The estimate of effective R is based on the combined analysis of deaths, case reporting, and hospitalizations where available. Current reported cases reflect infections 11-13 days prior, so estimates of effective R can only be made for the recent past.



**Figure 8.1: Percent of estimated COVID-19 infections detected.** This is estimated as the ratio of reported daily COVID-19 cases to estimated daily COVID-19 infections based on the SEIR disease transmission model. Due to measurement errors in cases and testing rates, the infection-detection rate can exceed 100% at particular points in time.



Estimated percent of circulating SARS-CoV-2 for primary variant families on December 12, 2022

Figure 9.1: Estimated percent of new infections that are Alpha variant




Figure 9.2: Estimated percent of new infections that are Beta variant



Figure 9.3: Estimated percent of new infections that are Delta variant




Figure 9.4: Estimated percent of new infections that are Gamma variant



Figure 9.5: Estimated percent of new infections that are BA.1/BA.2 variant






Figure 9.6: Estimated percent of new infections that are BA.5 variant



**Figure 10.1: Infection-fatality rate on December 12, 2022.** This is estimated as the ratio of COVID-19 deaths to estimated daily COVID-19 infections.



## Critical drivers

**Table 2: Current mandate implementation**

	Primary school closure	Secondary school closure	Higher school closure	Entry restrictions for some non-residents	Entry restrictions for all non-residents	Individual movements restricted	Curfew for businesses	Individual curfew	Gathering limit: 6 indoor, 10 outdoor	Gathering limit: 10 indoor, 25 outdoor	Gathering limit: 25 indoor, 50 outdoor	Gathering limit: 50 indoor, 100 outdoor	Gathering limit: 100 indoor, 250 outdoor	Restaurants closed	Bars closed	Restaurants / bars closed	Restaurants / bars curbside only	Gyms, pools, other leisure closed	Non-essential retail closed	Non-essential retail curbside only	Non-essential workplaces closed	Stay home order	Stay home fine	Mask mandate	Mask mandate fine	
Alabama																										
Alaska																										
Arizona																										
Arkansas																										
California																										
Colorado																										
Connecticut																										
Delaware																										
District of Columbia																										
Florida																										
Georgia																										
Hawaii																										
Idaho																										
Illinois																										
Indiana																										
Iowa																										
Kansas																										
Kentucky																										
Louisiana																										
Maine																										
Maryland																										
Massachusetts																										
Michigan																										
Minnesota																										
Mississippi																										
Missouri																										
Montana																										
Nebraska																										
Nevada																										
New Hampshire																										
New Jersey																										
New Mexico																										
New York																										
North Carolina																										
North Dakota																										
Ohio																										
Oklahoma																										
Oregon																										
Pennsylvania																										
Rhode Island																										
South Carolina																										
South Dakota																										
Tennessee																										
Texas																										
Utah																										
Vermont																										
Virginia																										
Washington																										
West Virginia																										
Wisconsin																										
Wyoming																										





Figure 11.1: Trend in the proportion of the population reporting always wearing a mask when leaving home





**Table 3: Estimates of vaccine effectiveness for specific vaccines used in the model at preventing severe disease and infection.** We use data from clinical trials directly, where available, and make estimates otherwise. More information can be found on our [website](#).

Vaccine	Effectiveness at preventing													
	Ancestral		Alpha		Beta		Gamma		Delta		BA.1/BA.2		BA.5	
	Severe disease	Infection	Severe disease	Infection	Severe disease	Infection	Severe disease	Infection	Severe disease	Infection	Severe disease	Infection	Severe disease	Infection
AstraZeneca	94%	63%	94%	63%	94%	69%	94%	69%	94%	69%	71%	36%	71%	36%
CanSino	66%	62%	66%	62%	64%	61%	64%	61%	64%	61%	48%	32%	48%	32%
CoronaVac	50%	47%	50%	47%	49%	46%	49%	46%	49%	46%	37%	24%	37%	24%
Covaxin	78%	73%	78%	73%	76%	72%	76%	72%	76%	72%	57%	38%	57%	38%
Johnson & Johnson	86%	72%	86%	72%	76%	64%	76%	64%	76%	64%	57%	33%	57%	33%
Moderna	97%	92%	97%	92%	97%	91%	97%	91%	97%	91%	73%	48%	73%	48%
Novavax	89%	83%	89%	83%	86%	82%	86%	82%	86%	82%	65%	43%	65%	43%
Pfizer/BioNTech	95%	86%	95%	86%	95%	84%	95%	84%	95%	84%	72%	44%	72%	44%
Sinopharm	73%	68%	73%	68%	71%	67%	71%	67%	71%	67%	53%	35%	53%	35%
Sputnik-V	92%	86%	92%	86%	89%	85%	89%	85%	89%	85%	67%	44%	67%	44%
Other vaccines	75%	70%	75%	70%	73%	69%	73%	69%	73%	69%	55%	36%	55%	36%
Other vaccines (mRNA)	91%	86%	91%	86%	88%	85%	88%	85%	88%	85%	67%	45%	67%	45%

Percent of the population having received at least one dose (12.1) and fully vaccinated against SARS-CoV-2 (12.2) by December 12, 2022

Figure 12.1: Percent of the population having received one dose of a COVID-19 vaccine




Figure 12.2: Percent of the population fully vaccinated against SARS-CoV-2







Figure 13.1: Estimated proportion of the total population that is not vaccinated but willing to be vaccinated as of June 24, 2022



**Figure 14.1: Percent of people who receive at least one dose of a COVID-19 vaccine and those who are fully vaccinated**



**Figure 15.1: Percent of people who are immune to Delta, BA.1/BA.2 or BA.5.** Immunity is based on protection due to prior vaccination and infection(s). Moreover, variant-specific immunity is also based on variant-variant specific protection.



## Projections and scenarios

Figure 16.1: Daily COVID-19 infections until April 01, 2023 for three scenarios




Figure 16.2: Daily COVID-19 reported cases until April 01, 2023 for three scenarios




Figure 16.3: Daily COVID-19 hospital census until April 01, 2023 for three scenarios




Figure 16.4: Reported daily COVID-19 deaths per 100,000






Figure 16.5: Total daily COVID-19 deaths per 100,000







**Figure 17.1: Comparison of reference model projections with other COVID modeling groups.** For this comparison, we are including projections of daily COVID-19 deaths from other modeling groups when available, last model update in brackets: the CDC Ensemble Model (CDC) [December 12, 2022]. Regional values are aggregates from available locations in that region.



**Figure 18.1: The estimated inpatient hospital usage is shown over time.** The percent of hospital beds occupied by COVID-19 patients is color-coded based on observed quantiles of the maximum proportion of beds occupied by COVID-19 patients. Less than 5% is considered *low stress*, 5-9% is considered *moderate stress*, 10-19% is considered *high stress*, and 20% or greater is considered *extreme stress*.



**Figure 19.1: The estimated intensive care unit (ICU) usage is shown over time.** The percent of ICU beds occupied by COVID-19 patients is color-coded based on observed quantiles of the maximum proportion of ICU beds occupied by COVID-19 patients. Less than 10% is considered *low stress*, 10-29% is considered *moderate stress*, 30-59% is considered *high stress*, and 60% or greater is considered *extreme stress*.



## More information

### Data sources:

Mask use and vaccine confidence data are from the [The Delphi Group at Carnegie Mellon University and University of Maryland COVID-19 Trends and Impact Surveys](#), in partnership with Facebook. Mask use data are also from [Premise](#), the Kaiser Family Foundation, and the [YouGov COVID-19 Behaviour Tracker](#) survey.

Genetic sequence and metadata are primarily from the GISAID Initiative. Further details available on the COVID-19 model [FAQ page](#).

### A note of thanks:

We wish to warmly acknowledge the support of [these](#) and others who have made our COVID-19 estimation efforts possible.

### More information:

For all COVID-19 resources at IHME, visit <http://www.healthdata.org/covid>.

To download our most recent results, visit our [Data downloads page](#).

Questions? Requests? Feedback? Please contact us at <https://www.healthdata.org/covid/contact-us>.