

Informe de resultados de COVID-19

Región de las Américas

08 de abril de 2021

Este documento contiene información resumida sobre las últimas proyecciones del modelo IHME sobre COVID-19 en la Región de las Américas. El modelo se ejecutó el 6 de abril de 2021 con datos hasta el 5 de abril de 2021.

Situación actual

- Los casos reportados diariamente en la última semana disminuyeron a 162,100 por día en promedio en comparación con 181,100 la semana anterior (Figura 1).
- Las muertes diarias en la última semana aumentaron a 5,600 por día en promedio en comparación con las 5,100 de la semana anterior (Figura 2). Esto convierte a COVID-19 en la principal causa de muerte en la Región de las Américas esta semana (Tabla 1).
- La tasa de mortalidad diaria es superior a 4 por millón en Argentina, Brasil, Chile, Ecuador, Paraguay, Perú y Uruguay (Figura 3).
- Estimamos que 28% de las personas en la Región de las Américas han sido infectadas al 5 de abril (Figura 4).
- La R efectiva, calculada usando casos, hospitalizaciones y muertes, es mayor que 1 en 78 países (Figura 5).
- La tasa de detección de infecciones en la Región de las Américas se acerca a 22% el 5 de abril (Figura 6).

Tendencias en los impulsores de la transmisión

- La movilidad la semana pasada fue 25% más baja que la línea de base anterior a COVID-19 (Figura 9). La movilidad estuvo cerca de la línea de base (dentro del 10%) en Colombia, El Salvador, Guatemala, Haití, Honduras y Nicaragua. La movilidad fue inferior a 30% de la línea de base en 12 países.
- Al 5 de abril, estimamos que 77% de las personas siempre usaban una máscara al salir de casa, sin cambios en comparación con la semana pasada (Figura 11). El uso de mascarillas fue inferior a 50% en 5 países.
- Hubo 177 pruebas de diagnóstico por cada 100,000 personas el 5 de abril (Figura 13).
- En la Región de las Américas, 79.9% de las personas dicen que aceptarían o probablemente aceptarían una vacuna para COVID-19. Esto es 1.5 puntos porcentuales más que la semana pasada. La fracción de la población que está dispuesta a recibir la vacuna COVID-19 oscila entre 38% en Jamaica 90% en Brasil (Figura 17).

- En nuestro escenario de referencia actual, esperamos que 1,029 millones de personas estén vacunados para el 1 de agosto (Figura 18).

Proyecciones

- En nuestro escenario de referencia, que representa lo que creemos que es más probable que suceda, nuestro modelo proyecta 1,993,000 muertes acumuladas el 1 de agosto de 2021. Esto representa 574,000 muertes adicionales del 5 de abril al 1 de agosto (Figura 19). Las muertes diarias alcanzarán un máximo de 7,050 el 24 de abril de 2021 (Figura 20).
- Si se alcanzara la cobertura universal de la mascarilla (95%) en la próxima semana, nuestro modelo proyecta 108.000 muertes menos en comparación con el escenario de referencia el 1 de agosto de 2021 (Figura 19).
- En nuestro peor escenario, nuestro modelo proyecta 2,261,000 muertes acumuladas el 1 de agosto de 2021, 268.000 muertes adicionales en comparación con nuestro escenario de referencia (Figura 19).
- Para el 1 de agosto, proyectamos que 286,700 vidas serán salvadas por el lanzamiento de la vacuna proyectada.
- La Figura 22 compara nuestros pronósticos de escenarios de referencia con otros modelos archivados públicamente. Los pronósticos son muy divergentes.
- En algún momento, desde abril hasta el 1 de agosto, 18 países tendrán una presión alta o extrema de camas de hospital (Figura 23). En algún momento, desde abril hasta el 1 de agosto, 18 países tendrán una presión alta o extrema en la capacidad de la UCI (Figura 24).

Actualizaciones de modelos

No hay actualizaciones importantes en el modelo esta semana.

COVID-19 Results Briefing

Region of the Americas

April 08, 2021

This document contains summary information on the latest projections from the IHME model on COVID-19 in the Region of the Americas. The model was run on April 6, 2021, with data through April 5, 2021.

Current situation

- Daily reported cases in the last week decreased to 162,100 per day on average compared to 181,100 the week before (Figure 1).
- Daily deaths in the last week increased to 5,600 per day on average compared to 5,100 the week before (Figure 2). This makes COVID-19 the number 1 cause of death in the Region of the Americas this week (Table 1).
- The daily death rate is greater than 4 per million in Argentina, Brazil, Chile, Ecuador, Paraguay, Peru, and Uruguay (Figure 3).
- We estimated that 28% of people in the Region of the Americas have been infected as of April 5 (Figure 4).
- Effective R, computed using cases, hospitalizations, and deaths, is greater than 1 in 78 countries (Figure 5).
- The infection detection rate in the Region of the Americas was close to 22% on April 5 (Figure 6).

Trends in drivers of transmission


- Mobility last week was 25% lower than the pre-COVID-19 baseline (Figure 9). Mobility was near baseline (within 10%) in Colombia, El Salvador, Guatemala, Haiti, Honduras, and Nicaragua. Mobility was lower than 30% of baseline in 12 countries.
- As of April 5, we estimated that 77% of people always wore a mask when leaving their home, unchanged from last week (Figure 11). Mask use was lower than 50% in five countries.
- There were 177 diagnostic tests per 100,000 people on April 5 (Figure 13).
- In the Region of the Americas, 79.9% of people say they would accept or would probably accept a vaccine for COVID-19. This is up by 1.5 percentage points from last week. The fraction of the population who are open to receiving a COVID-19 vaccine ranges from 38% in Jamaica to 90% in Brazil (Figure 17).
- In our current reference scenario, we expect that 1,029.19 million will be vaccinated by August 1 (Figure 18).

Projections

- In our **reference scenario**, which represents what we think is most likely to happen, our model projects 1,993,000 cumulative deaths on August 1, 2021. This represents 574,000 additional deaths from April 5 to August 1 (Figure 19). Daily deaths will peak at 7,050 on April 24, 2021 (Figure 20).
- If **universal mask coverage (95%)** were attained in the next week, our model projects 108,000 fewer cumulative deaths compared to the reference scenario on August 1, 2021 (Figure 19).
- Under our **worse scenario**, our model projects 2,261,000 cumulative deaths on August 1, 2021, an additional 268,000 deaths compared to our reference scenario (Figure 19).
- By August 1, we project that 286,700 lives will be saved by the projected vaccine rollout.
- Figure 22 compares our reference scenario forecasts to other publicly archived models. Forecasts are widely divergent.
- At some point from April through August 1, 18 countries will have high or extreme stress on hospital beds (Figure 23). At some point from April through August 1, 18 countries will have high or extreme stress on ICU capacity (Figure 24).

Model updates

There are no major updates in the model this week.

Figure 1. Reported daily COVID-19 cases

Table 1. Ranking of COVID-19 among the leading causes of mortality this week, assuming uniform deaths of non-COVID causes throughout the year

Cause name	Weekly deaths	Ranking
COVID-19	39,056	1
Ischemic heart disease	22,182	2
Stroke	10,124	3
Chronic obstructive pulmonary disease	7,401	4
Tracheal, bronchus, and lung cancer	6,369	5
Lower respiratory infections	6,211	6
Chronic kidney disease	6,184	7
Alzheimer's disease and other dementias	5,890	8
Diabetes mellitus	5,822	9
Cirrhosis and other chronic liver diseases	4,153	10

Figure 2. Reported daily COVID-19 deaths




Figure 3. Daily COVID-19 death rate per 1 million on April 05, 2021




Figure 4. Estimated percent of the population infected with COVID-19 on April 05, 2021




Figure 5. Mean effective R on March 25, 2021. The estimate of effective R is based on the combined analysis of deaths, case reporting, and hospitalizations where available. Current reported cases reflect infections 11-13 days prior, so estimates of effective R can only be made for the recent past. Effective R less than 1 means that transmission should decline, all other things being held the same.






Figure 6. Percent of COVID-19 infections detected. This is estimated as the ratio of reported daily COVID-19 cases to estimated daily COVID-19 infections based on the SEIR disease transmission model.



*Due to measurement errors in cases and testing rates, the infection to detection rate (IDR) can exceed 100% at particular points in time.

Figure 7. Percent of circulating SARS-CoV-2 for 3 primary variants on April 5, 2021.

A. Percent B.1.1.7 variant



B. Percent B.1.351 variant



C. Percent P1 variant





Figure 8. Infection fatality ratio on April 05, 2021. This is estimated as the ratio of COVID-19 deaths to infections based on the SEIR disease transmission model.



Critical drivers

Table 2. Current mandate implementation

	All nonessential businesses closed	Any businesses restricted	Any gatherings restricted	Mask use	School closure	Stay home order	Travel limits
Antigua and Barbuda	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Argentina	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Bahamas	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Barbados	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Belize	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Bolivia (Plurinational State of)	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Brazil	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Canada	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Chile	No mandate	Mandate in place	Mandate in place	No mandate	Mandate in place	Mandate in place	Mandate in place
Colombia	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Costa Rica	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Cuba	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Dominica	No mandate	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Dominican Republic	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Ecuador	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
El Salvador	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Grenada	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Guatemala	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Guyana	Mandate in place	Mandate in place	Mandate in place	No mandate	Mandate in place	Mandate in place	Mandate in place
Haiti	No mandate	Mandate in place	Mandate in place	Mandate in place	No mandate	No mandate	Mandate in place
Honduras	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Jamaica	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Mexico	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Nicaragua	No mandate	No mandate	No mandate	No mandate	Mandate in place	Mandate in place	Mandate in place
Panama	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Paraguay	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Peru	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Saint Kitts and Nevis	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Saint Lucia	Mandate in place	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Saint Vincent and the Grenadines	No mandate	Mandate in place	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Suriname	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Trinidad and Tobago	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
United States of America	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate
Uruguay	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place
Venezuela (Bolivarian Republic of)	No mandate	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place	Mandate in place





Figure 9. Trend in mobility as measured through smartphone app use compared to January 2020 baseline



— African Region — South-East Asia Region — Eastern Mediterranean Region
 — Region of the Americas — European Region — Western Pacific Region

Figure 10. Mobility level as measured through smartphone app use compared to January 2020 baseline (percent) on April 05, 2021




Figure 11. Trend in the proportion of the population reporting always wearing a mask when leaving home




Figure 12. Proportion of the population reporting always wearing a mask when leaving home on April 05, 2021




Figure 13. Trend in COVID-19 diagnostic tests per 100,000 people




Figure 14. COVID-19 diagnostic tests per 100,000 people on April 01, 2021




Figure 15. Increase in the risk of death due to pneumonia on February 1 2020 compared to August 1 2020




Table 3. The SEIR model uses variant-specific estimates of vaccine efficacy at preventing symptomatic disease and at preventing infection. We use data from clinical trials directly, where available, and make estimates otherwise. More information can be found on our website (<http://www.healthdata.org/node/8584>).

Vaccine	Efficacy at preventing disease: D614G & B.1.1.7	Efficacy at preventing infection: D614G & B.1.1.7	Efficacy at preventing disease: B.1.351 & P.1	Efficacy at preventing infection: B.1.351 & P.1
AstraZeneca	75%	52%	10%	7%
CanSinoBio	66%	57%	50%	44%
CoronaVac	50%	43%	38%	33%
Johnson & Johnson	72%	72%	64%	56%
Moderna	94%	85%	72%	62%
Novavax	89%	77%	49%	43%
Pfizer/BioNTech	91%	86%	69%	61%
Sinopharm	73%	63%	56%	48%
Sputnik V	92%	80%	70%	61%
Other mRNA vaccines	95%	83%	72%	63%
All other vaccines	75%	65%	57%	50%

Figure 16. Trend in the estimated proportion of the adult (18+) population that have been vaccinated or is open to receiving a COVID-19 vaccine based on Facebook survey responses (yes and yes, probably).




Figure 17. This figure shows the estimated proportion of the adult (18+) population that has been vaccinated or is open to receiving a COVID-19 vaccine based on Facebook survey responses (yes and yes, probably).





Figure 18. The number of people who receive any vaccine and those who are effectively vaccinated and protected against disease, accounting for efficacy, loss to follow up for two-dose vaccines, partial immunity after one dose, and immunity after two doses.



Projections and scenarios

We produce three scenarios when projecting COVID-19. The **reference scenario** is our forecast of what we think is most likely to happen:

- Vaccines are distributed at the expected pace.
- Governments adapt their response by re-imposing social distancing mandates for 6 weeks whenever daily deaths reach 8 per million, unless a location has already spent at least 7 of the last 14 days with daily deaths above this rate and not yet re-imposed social distancing mandates. In this case, the scenario assumes that mandates are re-imposed when daily deaths reach 15 per million.
- Variants B.1.1.7 (first identified in the UK), B.1.351 (first identified in South Africa), and P1 (first identified in Brazil) continue to spread from locations with (a) more than 5 sequenced variants, and (b) reports of community transmission, to adjacent locations following the speed of variant scale-up observed in the regions of the UK.
- In one-quarter of those vaccinated, mobility increases toward pre-COVID-19 levels.

The **worse scenario** modifies the reference scenario assumptions in three ways:

- First, it assumes that variants B.1.351 or P1 begin to spread within 3 weeks in adjacent locations that do not already have B.1.351 or P1 community transmission.
- Second, it assumes that all those vaccinated increase their mobility toward pre-COVID-19 levels.
- Third, it assumes that among those vaccinated, mask use starts to decline exponentially one month after completed vaccination.

The **universal masks scenario** makes all the same assumptions as the reference scenario but also assumes 95% of the population wear masks in public in every location.



Figure 19. Cumulative COVID-19 deaths until August 01, 2021 for three scenarios

Figure 20. Daily COVID-19 deaths until August 01, 2021 for three scenarios,


Figure 21. Daily COVID-19 infections until August 01, 2021 for three scenarios.




Figure 22. Comparison of reference model projections with other COVID modeling groups. For this comparison, we are including projections of daily COVID-19 deaths from other modeling groups when available: Delphi from the Massachusetts Institute of Technology (Delphi; <https://www.covidanalytics.io/home>), Imperial College London (Imperial; <https://www.covidsim.org>), The Los Alamos National Laboratory (LANL; <https://covid-19.bsvgateway.org/>), and the SI-KJalpha model from the University of Southern California (SIKJalpha; <https://github.com/scc-usc/ReCOVER-COVID-19>). Daily deaths from other modeling groups are smoothed to remove inconsistencies with rounding. Regional values are aggregates from available locations in that region.




Figure 23. The estimated inpatient hospital usage is shown over time. The percent of hospital beds occupied by COVID-19 patients is color coded based on observed quantiles of the maximum proportion of beds occupied by COVID-19 patients. Less than 5% is considered *low stress*, 5-9% is considered *moderate stress*, 10-19% is considered *high stress*, and greater than 20% is considered *extreme stress*.





Figure 24. The estimated intensive care unit (ICU) usage is shown over time. The percent of ICU beds occupied by COVID-19 patients is color coded based on observed quantiles of the maximum proportion of ICU beds occupied by COVID-19 patients. Less than 10% is considered *low stress*, 10-29% is considered *moderate stress*, 30-59% is considered *high stress*, and greater than 60% is considered *extreme stress*.



More information

Data sources:

Mask use data sources include [Premise](#); [Facebook Global Symptom Survey](#) (This research is based on survey results from University of Maryland Social Data Science Center) and the [Facebook United States Symptom Survey](#) (in collaboration with Carnegie Mellon University); Kaiser Family Foundation; [YouGov COVID-19 Behaviour Tracker](#) survey.

Vaccine hesitancy data are from the COVID-19 Beliefs, Behaviors, and Norms Study, a survey conducted on Facebook by the Massachusetts Institute of Technology (<https://covidsurvey.mit.edu/>).

Vaccine hesitancy data are from the [Facebook Global Symptom Survey](#) (This research is based on survey results from University of Maryland Social Data Science Center), the [Facebook United States Symptom Survey](#) (in collaboration with Carnegie Mellon University), and from the Facebook [COVID-19 Beliefs, Behaviors, and Norms Study](#) conducted by the Massachusetts Institute of Technology.

Genetic sequence and metadata are primarily from the GISAID Initiative. Further details available on the COVID-19 model [FAQ page](#).

A note of thanks:

We wish to warmly acknowledge the support of [these](#) and others who have made our COVID-19 estimation efforts possible.

More information:

For all COVID-19 resources at IHME, visit <http://www.healthdata.org/covid>.

Questions? Requests? Feedback? Please contact us at <https://www.healthdata.org/covid/contact-us>.