

COVID-19 Results Briefing

United Kingdom

November 18, 2021

This document contains summary information on the latest projections from the IHME model on COVID-19 in United Kingdom. The model was run on November 16, 2021, with data through November 15, 2021.

Current situation

- Daily infections in the last week increased to 60,600 per day on average compared to 55,400 the week before (Figure 1.1). Daily hospital census in the last week (through November 15) decreased to 8,800 per day on average compared to 9,200 the week before.
- Daily reported cases in the last week increased to 37,400 per day on average compared to 35,100 the week before (Figure 2.1).
- Reported deaths due to COVID-19 in the last week increased to 160 per day on average compared to 160 the week before (Figure 3.1).
- Total deaths due to COVID-19 in the last week increased to 180 per day on average compared to 180 the week before (Figure 3.1). This makes COVID-19 the number 2 cause of death in United Kingdom this week (Table 1). Estimated total daily deaths due to COVID-19 in the past week were 1.1 times larger than the reported number of deaths.
- No locations had daily reported COVID-19 death rates greater than 4 per million (Figure 4.1).
- No locations had daily total COVID-19 death rates greater than 4 per million (Figure 4.2).
- We estimate that 29% of people in United Kingdom have been infected as of November 15 (Figure 6.1).
- Effective R, computed using cases, hospitalizations, and deaths, is greater than 1 in 3 locations. (Figure 7.1).
- The infection-detection rate in United Kingdom was close to 74% on November 15 (Figure 8.1).
- Based on the GISAID and various national databases, combined with our variant spread model, we estimate the current prevalence of variants of concern (Figure 9.1). We estimate that the Beta variant is circulating in no countries, that the Delta variant is circulating in 4 countries, and that the Gamma variant is circulating in 1 country in the region.

Trends in drivers of transmission

- Mobility last week was 10% lower than the pre-COVID-19 baseline (Figure 11.1). Mobility was near baseline (within 10%) in England. Mobility was lower than 30% of baseline in no locations.
- As of November 15, in the COVID-19 Trends and Impact Survey, 27% of people self-report that they always wore a mask when leaving their home compared to 27% last week (Figure 13.1).
- There were 1266 diagnostic tests per 100,000 people on November 15 (Figure 15.1).
- As of November 15, 3 countries have reached 70% or more of the population who have received at least one vaccine dose and 2 countries have reached 70% or more of the population who are fully vaccinated (Figure 17.1).
- In our current reference scenario, we expect that 50.6 million people will be vaccinated with at least one dose by March 1 (Figure 20.1). We expect that 71% of the population will be fully vaccinated by March 1.
- Based on the estimate of the population that have been infected with COVID-19 and vaccinated to date, combined with assumptions on protection against infection with the Delta variant provided by either natural infection, vaccination or both, we estimate that 54% of the region is immune to the Delta variant. In our current reference scenario, we expect that by March 1, 69% of people will be immune to the Delta variant (Figure 21.1). These two calculations do not take into account waning of natural or vaccine-derived immunity.

Projections

- In our **reference scenario**, which represents what we think is most likely to happen, our model projects 195,000 cumulative reported deaths due to COVID-19 on March 1. This represents 28,000 additional deaths from November 15 to March 1. Daily reported deaths will rise to 340 by January 30, 2022 (Figure 22.1).
- Under our **reference scenario**, our model projects 222,000 cumulative total deaths due to COVID-19 on March 1. This represents 32,000 additional deaths from November 15 to March 1 (Figure 22.1).
- If **universal mask coverage (95%)** were attained in the next week, our model projects 17,000 fewer cumulative reported deaths compared to the reference scenario on March 1.
- Under our **worse scenario**, our model projects 215,000 cumulative reported deaths on March 1, an additional 20,000 deaths compared to our reference scenario. Daily reported deaths in the **worse scenario** will rise to 810 by January 28, 2022 (Figure 22.1).
- Daily infections in the **reference scenario** will rise to 119,410 by January 6, 2022 (Figure 22.3). Daily infections in the **worse scenario** will rise to 280,920 by January 5, 2022 (Figure 22.3).
- Daily cases in the **reference scenario** will rise to 86,330 by January 18, 2022 (Figure 22.4). Daily cases in the **worse scenario** will rise to 202,090 by January 16, 2022 (Figure 22.4).
- Daily hospital census in the **reference scenario** will rise to 18,160 by January 23, 2022 (Figure 22.5). Daily hospital census in the **worse scenario** will rise to 41,990 by January 21, 2022 (Figure 22.5).
- Figure 23.1 compares our reference scenario forecasts to other publicly archived models. Forecasts are widely divergent.
- At some point from November through March 1, 1 country will have high or extreme stress on hospital beds (Figure 24.1). At some point from November through March 1, 4 countries will have high or extreme stress on intensive care unit (ICU) capacity (Figure 25.1).

Model updates

No model updates.




Figure 1.1. Daily COVID-19 hospital census and infections

Figure 2.1. Reported daily COVID-19 cases, moving average


Table 1. Ranking of total deaths due to COVID-19 among the leading causes of mortality this week, assuming uniform deaths of non-COVID causes throughout the year

Cause name	Weekly deaths	Ranking
Ischemic heart disease	1,796	1
COVID-19	1,253	2
Stroke	974	3
Chronic obstructive pulmonary disease	845	4
Tracheal, bronchus, and lung cancer	824	5
Lower respiratory infections	805	6
Alzheimer's disease and other dementias	624	7
Colon and rectum cancer	466	8
Prostate cancer	307	9
Breast cancer	293	10

Figure 3.1. Smoothed trend estimate of reported daily COVID-19 deaths (blue) and total daily deaths due to COVID-19 (orange)



Daily COVID-19 death rate per 1 million on November 15, 2021

Figure 4.1 Daily reported COVID-19 death rate per 1 million





Figure 4.2 Daily total COVID-19 death rate per 1 million



Cumulative COVID-19 deaths per 100,000 on November 15, 2021

Figure 5.1 Reported cumulative COVID-19 deaths per 100,000




Figure 5.2 Total cumulative COVID-19 deaths per 100,000




Figure 6.1. Estimated percent of the population infected with COVID-19 on November 15, 2021




Figure 7.1. Mean effective R on November 4, 2021. Effective R less than 1 means that transmission should decline, all other things being held the same. The estimate of effective R is based on the combined analysis of deaths, case reporting, and hospitalizations where available. Current reported cases reflect infections 11-13 days prior, so estimates of effective R can only be made for the recent past.





Figure 8.1. Percent of COVID-19 infections detected. This is estimated as the ratio of reported daily COVID-19 cases to estimated daily COVID-19 infections based on the SEIR disease transmission model. Due to measurement errors in cases and testing rates, the infection-detection rate can exceed 100% at particular points in time.



Estimated percent of circulating SARS-CoV-2 for primary variant families on November 15, 2021

Figure 9.1 Estimated percent Alpha variant




Figure 9.2 Estimated percent Beta variant




Figure 9.3 Estimated percent Delta variant




Figure 9.4 Estimated percent Gamma variant





Figure 10.1. Infection-fatality rate on November 15, 2021. This is estimated as the ratio of COVID-19 deaths to estimated daily COVID-19 infections.



Critical drivers

Table 2. Current mandate implementation

	Primary school closure	Secondary school closure	Higher school closure	Borders closed to any non-resident	Borders closed to all non-residents	Individual movements restricted	Curfew for businesses	Individual curfew	Gathering limit: 6 indoor, 10 outdoor	Gathering limit: 10 indoor, 25 outdoor	Gathering limit: 25 indoor, 50 outdoor	Gathering limit: 50 indoor, 100 outdoor	Gathering limit: 100 indoor, 250 outdoor	Restaurants closed	Bars closed	Restaurants / bars closed	Restaurants / bars curbside only	Gyms, pools, other leisure closed	Non-essential retail closed	Non-essential retail curbside only	Non-essential workplaces closed	Stay home order	Stay home fine	Mask mandate	Mask mandate fine
England	No mandate	No mandate	Mandate in place (updated from previous reporting)	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate
Northern Ireland	No mandate	No mandate	Mandate in place (imposed this week)	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	Mandate in place (imposed this week)	Mandate in place (imposed this week)	Mandate in place (imposed this week)	Mandate in place (imposed this week)	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	Mandate in place (updated from previous reporting)
Scotland	No mandate	No mandate	Mandate in place (updated from previous reporting)	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	Mandate in place (updated from previous reporting)
Wales	No mandate	No mandate	Mandate in place (updated from previous reporting)	No mandate	No mandate	Mandate in place (updated from previous reporting)	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	No mandate	Mandate in place (updated from previous reporting)

Figure 11.1. Trend in mobility as measured through smartphone app use, compared to January 2020 baseline




Figure 12.1. Mobility level as measured through smartphone app use, compared to January 2020 baseline (percent) on November 15, 2021




Figure 13.1. Trend in the proportion of the population reporting always wearing a mask when leaving home




Figure 14.1. Proportion of the population reporting always wearing a mask when leaving home on November 15, 2021




Figure 15.1. Trend in COVID-19 diagnostic tests per 100,000 people




Figure 16.1. COVID-19 diagnostic tests per 100,000 people on November 15, 2021




Table 3. Estimates of vaccine efficacy for specific vaccines used in the model at preventing disease and infection. The SEIR model uses variant-specific estimates of vaccine efficacy at preventing symptomatic disease and at preventing infection. We use data from clinical trials directly, where available, and make estimates otherwise. More information can be found on our [website](#).

Vaccine	Efficacy at preventing disease: ancestral and Alpha	Efficacy at preventing infection: ancestral and Alpha	Efficacy at preventing disease: Beta, Delta, & Gamma	Efficacy at preventing infection: Beta, Delta, & Gamma
AstraZeneca	90%	52%	85%	49%
CoronaVac	50%	44%	43%	38%
Covaxin	78%	69%	68%	60%
Johnson & Johnson	86%	72%	60%	56%
Moderna	94%	89%	94%	80%
Novavax	89%	79%	79%	69%
Pfizer/BioNTech	94%	86%	85%	78%
Sinopharm	73%	65%	63%	56%
Sputnik-V	92%	81%	80%	70%
Tianjin	66%	58%	57%	50%
CanSino				
Other vaccines	75%	66%	65%	57%
Other vaccines (mRNA)	91%	86%	85%	78%

Percent of the population having received at least one dose (17.1) and fully vaccinated against SARS-CoV-2 (17.2) by November 15, 2021

Figure 17.1 Percent of the population having received one dose of a COVID-19 vaccine



Figure 17.2 Percent of the population fully vaccinated against SARS-CoV-2



Figure 20.1. Percent of people who receive at least one dose of a COVID-19 vaccine and those who are fully vaccinated





Figure 21.1. Percentage of people who are immune to non-escape variants and the percentage of people who are immune to escape variants



Projections and scenarios

We produce three scenarios when projecting COVID-19. The **reference scenario** is our forecast of what we think is most likely to happen:

- Vaccines are distributed at the expected pace. Brand- and variant-specific vaccine efficacy is updated using the latest available information from peer-reviewed publications and other reports.
- Future mask use is the mean of mask use over the last 7 days.
- Mobility increases as vaccine coverage increases.
- Governments adapt their response by re-imposing social distancing mandates for 6 weeks whenever daily deaths reach 8 per million, unless a location has already spent at least 7 of the last 14 days with daily deaths above this rate, and not yet re-imposed social distancing mandates. In this case, the reference scenario assumes that mandates are re-imposed when daily deaths reach 15 per million.
- Variants Alpha, Beta, Gamma, and Delta continue to spread regionally and globally from locations with sufficient transmission.

The **worse scenario** modifies the reference scenario assumption in four ways:

- 100% of vaccinated individuals stop using masks.
- Mobility increases in all locations to 25% above the pre-pandemic winter baseline, irrespective of vaccine coverage.
- Governments are more reluctant to re-impose social distancing mandates, waiting until the daily death rate reaches 15 per million, unless a location has already spent at least 7 of the last 14 days with daily deaths above this rate, and not yet re-imposed social distancing mandates. In this case, the reference scenario assumes that mandates are re-imposed when daily deaths reach 38 per million. In either case, we assume social distancing mandates remain in effect for 6 weeks.
- Variants Alpha, Beta, Gamma, and Delta spread between locations twice as fast when compared with our reference scenario.

The **universal masks scenario** makes all the same assumptions as the reference scenario but assumes all locations reach 95% mask use within 7 days.

Daily COVID-19 deaths until March 01, 2022 for three scenarios

Figure 22.1 Reported daily COVID-19 deaths per 100,000




Figure 22.2 Total daily COVID-19 deaths per 100,000






Figure 22.3. Daily COVID-19 infections until March 01, 2022 for three scenarios

Figure 22.4. Daily COVID-19 reported cases until March 01, 2022 for three scenarios


Figure 22.5. Daily COVID-19 hospital census until March 01, 2022 for three scenarios




Figure 23.1. Comparison of reference model projections with other COVID modeling groups. For this comparison, we are including projections of daily COVID-19 deaths from other modeling groups when available, last model update in brackets: Delphi from the Massachusetts Institute of Technology ([Delphi](#)) [November 17, 2021], Imperial College London ([Imperial](#)) [November 3, 2021], the SI-KJalpha model from the University of Southern California ([SIKJalpha](#)) [November 17, 2021]. Daily deaths from other modeling groups are smoothed to remove inconsistencies with rounding. Regional values are aggregates from available locations in that region.




Figure 24.1. The estimated inpatient hospital usage is shown over time. The percent of hospital beds occupied by COVID-19 patients is color-coded based on observed quantiles of the maximum proportion of beds occupied by COVID-19 patients. Less than 5% is considered *low stress*, 5-9% is considered *moderate stress*, 10-19% is considered *high stress*, and 20% or greater is considered *extreme stress*.





Figure 25.1. The estimated intensive care unit (ICU) usage is shown over time. The percent of ICU beds occupied by COVID-19 patients is color-coded based on observed quantiles of the maximum proportion of ICU beds occupied by COVID-19 patients. Less than 10% is considered *low stress*, 10-29% is considered *moderate stress*, 30-59% is considered *high stress*, and 60% or greater is considered *extreme stress*.



More information

Data sources:

Mask use and vaccine confidence data are from the [The Delphi Group at Carnegie Mellon University and University of Maryland COVID-19 Trends and Impact Surveys](#), in partnership with Facebook. Mask use data are also from [Premise](#), the Kaiser Family Foundation, and the [YouGov COVID-19 Behaviour Tracker](#) survey.

Genetic sequence and metadata are primarily from the GISAID Initiative. Further details available on the COVID-19 model [FAQ page](#).

A note of thanks:

We wish to warmly acknowledge the support of [these](#) and others who have made our COVID-19 estimation efforts possible.

More information:

For all COVID-19 resources at IHME, visit <http://www.healthdata.org/covid>.

To download our most recent results, visit our [Data downloads page](#).

Questions? Requests? Feedback? Please contact us at <https://www.healthdata.org/covid/contact-us>.