

Risk Factor Definitions

Hypertension – Treatment	Among adults age 30 and older who have been previously diagnosed and/or have systolic blood pressure 140 mmHg or higher, the proportion who are receiving treatment.
Hypertension – Control	Among adults age 30 and older who have been previously diagnosed and/or have systolic blood pressure 140 mmHg or higher, the proportion who are receiving treatment and currently have systolic blood pressure less than 140 mmHg.
Alcohol use – Any drinking	Among all adults age 21 and older, the proportion who have had one or more alcoholic drinks in the previous month.
Alcohol use – Binge drinking	Among all adults age 21 and older, the proportion who have had at least four (for women) or five (for men) alcoholic drinks in a single occasion one or more times in the previous month.
Alcohol use – Heavy drinking	Among all adults age 21 and older, the proportion who have had, on average, more than one (for women) or two (for men) alcoholic drinks per day during the previous month.
Alcohol use – Binge drinking among drinkers	Among adults age 21 and older who have had one or more alcoholic drinks in the previous month, the proportion who have had at least four (for women) or five (for men) alcoholic drinks in a single occasion one or more times in the previous month.
Alcohol use – Heavy drinking among drinkers	Among adults age 21 and older who have had one or more alcoholic drinks in the previous month, the proportion who have had, on average, more than one (for women) or two (for men) alcoholic drinks per day during the previous month.