

ANNEX

METHODS


The analytical strategy of GBD

The GBD approach contains 18 distinct components, as outlined in Figure A1. The components of GBD are interconnected. For example, when new data is incorporated into the age-specific mortality rates analysis (component 2), other dependent components must also be updated, such as rescaling deaths for each cause (component 5), healthy life expectancy or HALE (component 12), YLLs (component 13), and estimation of YLLs attributable to each risk factor (component 18). The inner workings of key components are briefly described in this publication, and more detailed descriptions of each component are included in the published articles.

Estimating age- and sex-specific mortality

Researchers identified sources of under-5 and adult mortality data from vital and sample registration systems as well as from surveys that ask mothers about live births and deaths of their children and ask people about siblings and their survival. Researchers processed that data to address biases and estimated the probability of death between ages 0 and 5 and ages 15 and 60 using statistical models. Finally, researchers used these probability estimates as well as a model life table system to estimate age-specific mortality rates by sex between 1970 and 2010.

Figure A1: The 18 components of GBD and their interrelations




Estimating years lost due to premature death

Researchers compiled all available data on causes of death from 187 countries. Information about causes of death was derived from vital registration systems, mortality surveillance systems, censuses, surveys, hospital records, police records, mortuaries, and verbal autopsies. Verbal autopsies are surveys that collect information from individuals familiar with the deceased about the signs and symptoms the person had prior to death. GBD 2010 researchers closely examined the completeness of the data. For those countries where cause of death data were incomplete, researchers used statistical techniques to compensate for the inherent biases. They also standardized causes of death across different data sources by mapping different versions of the International Classification of Diseases coding system to the GBD cause list.

Next, researchers examined the accuracy of the data, scouring rows and rows of data for “garbage codes.” Garbage codes are misclassifications of death in the data, and researchers identified thousands of them. Some garbage codes are instances where we know the cause listed cannot possibly lead to death. Examples found in records include “abdominal rigidity,” “senility,” and “yellow nail syndrome.” To correct these, researchers drew on evidence from medical literature, expert judgment, and statistical techniques to reassign each of these to more probable causes of death.

After addressing data-quality issues, researchers used a variety of statistical models to determine the number of deaths from each cause. This approach, named Cause of Death Ensemble modeling or CODEm, was designed based on statistical tech-

Figure A2: Leading causes of death and premature death in sub-Saharan Africa, 2010



niques called “ensemble modeling.” Ensemble modeling was made famous by the recipients of the Netflix Prize in 2009, BellKor’s Pragmatic Chaos, who engineered the best algorithm to predict how much a person would like a film, taking into account their movie preferences.

To ensure that the number of deaths from each of cause did not exceed the total number of deaths estimated in a separate GBD demographic analysis, researchers applied a correction technique named CoDCorrect. This technique makes certain that estimates of the number of deaths from each cause do not add up to more than 100% of deaths in a given year.

After producing estimates of the number of deaths from each of the 235 fatal outcomes included in the GBD cause list, researchers then calculated years of life lost to premature death, or YLLs. For every death from a particular cause, researchers estimated the number of years lost based on the highest life expectancy in the deceased’s age group. For example, if a 20-year-old male died in a car accident in South Africa in 2010, he has 66 years of life lost, that is, the highest remaining life expectancy in 20-year-olds, as experienced by 20-year-old females in Japan.

When comparing rankings of the leading causes of death versus YLLs, YLLs place more weight on the causes of death that occur in younger age groups, as shown in Figure A2. For example, malaria represents a greater percentage of total YLLs than total deaths since it is a leading killer of children under age 5. Ischemic heart disease, by contrast, accounts for a smaller percentage of total YLLs than total deaths, as it primarily kills older people.

Estimating years lived with disability

Researchers estimated the prevalence of each sequela using different sources of data, including government reports of cases of infectious diseases, data from population-based disease registries for conditions such as cancers and chronic kidney diseases, antenatal clinic data, hospital discharge data, data from outpatient facilities, interview questions, and direct measurements of hearing, vision, and lung function testing from surveys and other sources.

Confronted with the challenge of data gaps in many regions and for numerous types of sequelae, they developed a statistical modeling tool named DisMod-MR (for Disease Modeling – Metaregression) to estimate prevalence using available data on incidence, prevalence, remission, duration, and extra risk of mortality due to the disease.

Researchers estimated disability weights using data collected from almost 14,000 respondents via household surveys in Bangladesh, Indonesia, Peru, Tanzania, and the United States. Disability weights measure the severity of different sequelae that result from disease and injury. Data were also used from an Internet survey of more than 16,000 people. GBD researchers presented different lay definitions of sequelae grouped into 220 unique health states to survey respondents, and respondents were then asked to rate the severity of the different health states. The results were similar

across all surveys despite cultural and socioeconomic differences. Respondents consistently placed health states such as mild hearing loss and long-term treated fractures at the low end of the severity scale, while they ranked acute schizophrenia and severe multiple sclerosis as very severe.

Finally, years lived with disability, or YLDs, are calculated as prevalence of a sequela multiplied by the disability weight for that sequela. The number of years lived with disability for a specific disease or injury are calculated as the sum of the YLDs from each sequela arising from that cause.

Estimating disability-adjusted life years

Disability-adjusted life years (DALYs) were calculated by adding together YLLs and YLDs. Figure A3 compares the 10 leading diseases and injuries calculated as percentages of both regional deaths and regional DALYs. This figure also shows the top 10 risk factors attributable to deaths and DALYs in the sub-Saharan Africa region. DALYs are a powerful tool for priority setting as they measure disease burden from non-fatal, as well as fatal, conditions. Yet another reason why top causes of DALYs differ from leading causes of death is that DALYs give more weight to death in younger ages, as illustrated by the case of neonatal encephalopathy. In contrast, stroke causes a larger percentage of total deaths than DALYs, as it primarily impacts older people.

Estimating DALYs attributable to risk factors


To estimate the number of healthy years lost, or DALYs, attributable to potentially avoidable risk factors, researchers collected detailed data on exposure to different risk factors. The study used data from sources such as satellite data on air pollution, breastfeeding data from population surveys, and blood and bone lead levels from medical examination surveys and epidemiological surveys. Researchers then collected data on the effects of risk factors on disease outcomes through systematic reviews of epidemiological studies.

All risk factors analyzed met common criteria in four areas:

1. The likely importance of a risk factor for policymaking or disease burden.
2. Availability of sufficient data to estimate exposure to a particular risk factor.
3. Rigorous scientific evidence that specific risk factors cause certain diseases and injuries.
4. Scientific findings about the effects of different risk factors that are relevant for the general population.

To calculate the number of DALYs attributable to different risk factors, researchers compared the disease burden in a group exposed to a risk factor to the disease burden in a group that had zero exposure to that risk factor. When subjects with zero exposure were impossible to find, as in the case of high blood pressure, for example, researchers established a level of minimum exposure that leads to the best health outcomes.

Figure A3: The 10 leading diseases and injuries and 10 leading risk factors based on percentage of deaths and DALYs in sub-Saharan Africa, 2010



Note: This figure compares the percent of DALYs and deaths attributable to different diseases and injuries (shown in blue) as well as risk factors (shown in red). Certain causes, such as iron deficiency, cause more DALYs than they do deaths. DALYs are an important tool for decision-makers because they capture years of healthy life lost from both premature death and years lived with disability.

Table A1: Age-standardized death rates, years of life lost, and years lived with disability, and life expectancy at birth and healthy life expectancy at birth for 1990 and 2010 for both sexes combined


Country	Age-standardized death rate (per 100,000)				Age-standardized YLL rate (per 100,000)			
	1990		2010		1990		2010	
	Rate	Rank	Rate	Rank	Rate	Rank	Rate	Rank
Angola	1,995	43	1,257	20	82,686	42	44,380	14
	(1,527-2,733)	(29-46)	(993-1,644)	(8-40)	(66,351-109,548)	(30-46)	(35,634-57,681)	(11-36)
Benin	1,312	19	1,080	11	58,713	24	38,726	13
	(1,244-1,388)	(13-22)	(1,014-1,151)	(8-14)	(54,284-63,355)	(21-26)	(35,694-42,020)	(11-17)
Botswana	951	28	715	2	31,563	4	26,692	4
	(848-1,057)	(2-4)	(599-821)	(2-4)	(28,522-34,692)	(3-5)	(23,854-29,414)	(3-5)
Burkina Faso	1,521	28	1,396	33	68,852	30	59,507	39
	(1,459-1,585)	(25-31)	(1,288-1,503)	(26-37)	(65,046-73,156)	(27-34)	(52,426-67,312)	(31-42)
Burundi	1,966	42	1,759	40	80,347	39	62,752	40
	(1,517-2,692)	(30-46)	(1,332-2,427)	(31-46)	(68,981-106,923)	(30-46)	(49,931-85,236)	(27-44)
Cameroon	1,296	16	1,277	23	51,072	17	49,262	26
	(1,243-1,345)	(13-20)	(1,178-1,377)	(18-30)	(48,257-53,902)	(16-20)	(44,889-53,932)	(21-31)
Cape Verde	785	1	586	1	25,843	2	16,730	1
	(746-827)	(1-1)	(497-685)	(1-1)	(24,238-27,543)	(1-2)	(13,743-20,284)	(1-2)
Central African Republic	2,131	45	2,408	46	81,026	44	90,581	44
	(2,010-2,256)	(43-46)	(2,138-2,713)	(44-46)	(76,404-86,270)	(39-45)	(80,204-101,333)	(44-46)
Chad	1,508	28	1,389	32	67,610	29	58,839	37
	(1,426-1,596)	(24-31)	(1,284-1,514)	(24-37)	(62,369-73,496)	(26-34)	(52,545-66,878)	(31-42)
Comoros	1,439	24	1,223	18	52,340	19	38,959	14
	(1,308-1,590)	(19-30)	(1,092-1,373)	(13-29)	(48,247-57,207)	(16-22)	(34,801-44,336)	(9-18)
Congo	1,696	35	1,401	34	60,758	35	49,063	25
	(1,565-1,848)	(30-42)	(1,281-1,540)	(26-37)	(55,481-66,618)	(21-29)	(43,995-54,467)	(20-32)
Côte d'Ivoire	1,400	25	1,457	36	58,331	23	56,879	35
	(1,339-1,466)	(21-25)	(1,341-1,576)	(31-38)	(55,059-61,623)	(21-26)	(51,282-62,523)	(31-39)
Democratic Republic of the Congo	1,568	29	1,506	37	65,799	27	59,031	38
	(1,467-1,663)	(26-33)	(1,359-1,615)	(33-39)	(61,413-70,116)	(26-31)	(53,984-63,582)	(33-40)
Eritrea	1,807	41	1,353	30	67,111	28	44,346	19
	(1,694-1,922)	(36-43)	(1,228-1,466)	(20-36)	(62,000-72,863)	(27-33)	(39,865-48,958)	(15-26)
Ethiopia	2,152	46	1,292	25	86,420	45	43,719	18
	(2,044-2,278)	(43-46)	(1,232-1,352)	(19-30)	(80,270-93,661)	(42-46)	(41,409-46,267)	(15-22)
Gabon	1,212	18	1,432	35	46,803	15	49,393	27
	(1,218-1,412)	(11-23)	(1,267-1,637)	(25-40)	(42,373-50,001)	(12-16)	(42,613-58,336)	(19-37)
Gambia, The	1,315	17	1,109	12	54,039	20	41,302	15
	(1,146-1,518)	(9-27)	(900-1,356)	(6-27)	(48,109-60,649)	(16-25)	(34,115-49,904)	(8-26)
Ghana	1,185	10	1,030	8	45,628	13	35,128	8
	(1,115-1,253)	(8-12)	(968-1,086)	(6-12)	(42,476-49,061)	(12-15)	(32,885-37,437)	(7-12)
Guinea	1,508	26	1,233	19	69,094	31	48,350	23
	(1,444-1,580)	(24-30)	(1,152-1,324)	(15-27)	(64,734-74,410)	(27-34)	(43,740-53,673)	(18-31)
Guinea-Bissau	1,603	31	1,391	31	72,398	33	55,214	33
	(1,407-1,831)	(24-41)	(1,153-1,649)	(17-39)	(64,267-80,951)	(27-41)	(46,304-65,422)	(24-40)
Kenya	1,092	5	1,013	7	39,731	9	36,153	11
	(1,041-1,141)	(4-8)	(940-1,081)	(6-11)	(37,585-42,064)	(6-11)	(33,852-38,494)	(8-13)
Lesotho	1,316	20	2,130	44	43,569	12	85,888	45
	(1,245-1,394)	(13-23)	(1,930-2,341)	(43-46)	(41,196-46,177)	(11-15)	(78,939-92,875)	(44-46)
Liberia	1,585	30	1,333	28	75,995	37	53,808	32
	(1,482-1,728)	(26-36)	(1,244-1,425)	(22-34)	(70,210-82,809)	(33-43)	(49,728-57,821)	(28-36)
Madagascar	1,451	25	1,156	14	56,586	22	37,200	12
	(1,391-1,506)	(23-27)	(1,062-1,255)	(11-21)	(53,790-59,954)	(20-24)	(34,362-40,628)	(8-15)
Malawi	1,768	40	1,730	42	79,007	43	66,486	42
	(1,685-1,843)	(35-43)	(1,591-1,843)	(39-43)	(74,928-83,200)	(36-44)	(61,969-70,511)	(40-43)
Mali	1,726	38	1,331	27	78,963	41	53,703	31
	(1,661-1,800)	(33-41)	(1,241-1,425)	(20-34)	(74,141-84,559)	(36-44)	(48,704-59,440)	(27-37)
Mauritania	1,244	12	1,031	9	45,799	14	36,003	10
	(1,194-1,298)	(10-17)	(955-1,110)	(6-12)	(43,134-48,769)	(12-15)	(32,543-39,793)	(7-14)
Mauritius	1,014	3	766	3	24,451	1	17,273	2
	(1,000-1,024)	(2-4)	(751-776)	(2-5)	(24,031-24,836)	(1-2)	(16,915-17,579)	(1-2)
Mozambique	1,659	32	1,725	41	74,370	35	67,547	43
	(1,585-1,735)	(30-37)	(1,577-1,852)	(39-43)	(69,485-79,516)	(32-41)	(62,839-72,350)	(39-43)
Namibia	1,259	13	1,298	26	39,681	8	42,112	16
	(1,215-1,306)	(11-18)	(1,187-1,418)	(18-32)	(38,003-41,586)	(6-10)	(37,956-46,928)	(13-21)
Niger	1,765	39	1,268	22	88,497	46	52,955	30
	(1,673-1,866)	(34-42)	(1,184-1,369)	(16-31)	(82,055-95,891)	(44-46)	(47,331-59,543)	(23-38)
Nigeria	1,343	21	1,159	15	60,604	26	49,276	28
	(1,267-1,414)	(16-23)	(1,084-1,237)	(12-19)	(56,578-64,799)	(23-27)	(45,962-52,718)	(22-30)
Rwanda	2,000	44	1,062	10	76,083	38	35,591	9
	(1,874-2,118)	(41-45)	(994-1,129)	(7-13)	(71,263-80,773)	(33-42)	(33,361-37,861)	(7-13)
São Tomé and Príncipe	1,044	4	794	4	39,144	6	24,592	3
	(985-1,099)	(3-5)	(726-874)	(2-5)	(36,220-42,042)	(6-11)	(22,306-27,110)	(3-4)
Senegal	1,278	14	1,009	6	49,632	16	33,824	7
	(1,235-1,331)	(12-19)	(947-1,079)	(6-11)	(47,252-52,338)	(15-19)	(31,796-35,962)	(7-10)
Seychelles	1,133	7	1,187	16	29,858	3	31,195	6
	(1,099-1,162)	(5-10)	(1,116-1,238)	(12-21)	(28,919-30,862)	(3-4)	(29,079-32,715)	(5-7)
Sierra Leone	1,682	34	1,347	29	78,271	40	49,641	29
	(1,578-1,773)	(31-40)	(1,263-1,423)	(23-34)	(72,119-83,863)	(34-44)	(45,708-53,125)	(22-31)
Somalia	1,738	36	1,606	39	70,184	32	57,208	34
	(1,407-2,201)	(24-45)	(1,276-2,027)	(24-43)	(59,274-86,060)	(24-44)	(46,126-70,933)	(22-43)
South Africa	1,133	8	1,266	21	34,540	5	48,286	24
	(1,071-1,180)	(5-10)	(1,200-1,334)	(17-29)	(32,586-36,273)	(4-5)	(45,928-51,074)	(20-30)
Sudan	1,110	6	799	5	41,735	11	28,295	5
	(1,054-1,165)	(5-9)	(744-860)	(3-5)	(39,218-44,421)	(8-12)	(26,208-30,917)	(4-6)
Swaziland	1,239	11	2,186	45	39,973	10	80,065	44
	(1,163-1,316)	(10-18)	(1,950-2,429)	(43-46)	(37,549-42,545)	(6-11)	(72,966-87,908)	(43-45)
Tanzania	1,357	22	1,137	13	55,603	21	43,461	17
	(1,305-1,407)	(18-23)	(1,053-1,223)	(11-18)	(52,816-58,575)	(19-24)	(40,811-46,456)	(15-21)
Togo	1,282	15	1,199	17	51,184	18	46,594	22
	(1,229-1,337)	(12-19)	(1,106-1,296)	(13-23)	(48,232-54,390)	(16-20)	(42,401-51,158)	(17-28)
Uganda	1,658	33	1,290	24	75,253	36	45,587	21
	(1,563-1,760)	(30-38)	(1,196-1,387)	(18-31)	(71,264-79,804)	(33-41)	(42,676-48,830)	(17-25)
Zambia	1,722	37	1,533	38	73,053	34	57,620	36
	(1,647-1,792)	(33-41)	(1,414-1,644)	(33-40)	(69,561-76,818)	(30-39)	(53,531-61,944)	(31-39)
Zimbabwe	1,144	9	1,801	43	39,368	7	65,919	41
	(1,079-1,201)	(5-10)	(1,601-2,023)	(40-43)	(37,293-41,290)	(6-10)	(59,728-72,926)	(41-43)

Age-standardized YLD rate (per 100,000)				Life expectancy at birth				Health-adjusted life expectancy at birth			
1990		2010		1990		2010		1990		2010	
Rate	Rank	Rate	Rank	LE	Rank	LE	Rank	HALE	Rank	HALE	Rank
15,039	30	13,712	12	47.9	41	61.0	19	40.5	40	52.0	19
(12,384-18,328)	(13-46)	(11,204-16,796)	(4-38)	(41.8-53.1)	(28-46)	(54.3-67.0)	(6-39)	(35.4-44.8)	(28-46)	(46.5-56.7)	(7-38)
15,237	37	13,830	17	55.9	23	63.4	13	46.6	24	53.7	14
(12,408-18,418)	(17-46)	(11,270-16,742)	(6-39)	(55.1-56.8)	(20-26)	(60.8-65.7)	(8-20)	(44.6-46.4)	(21-26)	(50.9-56.4)	(8-21)
13,528	8	14,557	34	66.7	3	71.2	3	56.6	4	59.3	4
(11,131-16,243)	(3-28)	(12,007-17,522)	(10-45)	(64.2-69.4)	(2-5)	(67.7-75.4)	(1-5)	(53.9-59.4)	(3-5)	(55.9-63.1)	(3-6)
14,862	29	13,409	11	52.1	30	55.3	38	43.8	30	47.2	37
(12,335-17,843)	(14-44)	(10,997-16,422)	(4-30)	(51.2-53.1)	(27-34)	(49.9-59.6)	(26-44)	(42.0-45.5)	(27-34)	(42.3-51.2)	(23-43)
14,581	23	14,412	32	48.9	39	54.6	39	41.6	37	46.4	39
(12,072-17,535)	(9-42)	(11,821-17,419)	(11-44)	(41.7-54.7)	(25-46)	(46.4-62.1)	(17-46)	(36.1-48.4)	(26-46)	(39.8-52.5)	(18-46)
14,617	25	13,891	19	58.3	19	59.1	26	49.1	18	50.2	26
(12,034-17,487)	(9-43)	(11,399-16,769)	(7-39)	(57.6-59.1)	(16-20)	(56.1-61.9)	(17-36)	(47.2-50.8)	(16-20)	(47.2-52.8)	(17-35)
13,500	6	12,947	6	69.8	1	75.1	1	59.1	2	63.8	2
(10,843-18,557)	(3-46)	(10,310-20,228)	(2-46)	(68.8-70.9)	(1-2)	(71.7-78.2)	(1-3)	(55.2-61.4)	(1-4)	(58.3-67.3)	(1-4)
15,696	46	15,022	40	48.4	44	46.4	46	40.8	44	39.7	46
(12,822-18,800)	(22-46)	(12,185-18,088)	(23-46)	(46.9-49.8)	(37-44)	(41.9-50.7)	(43-46)	(38.9-42.6)	(36-45)	(35.7-43.3)	(43-46)
15,080	34	14,364	30	52.5	28	55.6	29	43.9	29	46.9	38
(12,463-18,027)	(17-45)	(11,770-17,273)	(14-42)	(51.6-53.3)	(27-33)	(50.3-59.8)	(24-43)	(42.1-45.6)	(27-34)	(42.3-50.7)	(25-44)
13,050	4	12,879	19	57.6	19	62.8	14	49.6	17	54.1	12
(10,566-15,698)	(3-16)	(10,556-15,420)	(3-20)	(54.5-60.3)	(14-26)	(59.2-66.2)	(7-27)	(46.7-52.1)	(12-22)	(50.6-57.4)	(7-23)
15,443	42	14,357	31	48.9	26	58.9	27	46.0	26	50.0	27
(12,837-18,350)	(21-46)	(11,832-17,146)	(14-42)	(52.8-56.6)	(21-29)	(55.2-61.9)	(16-37)	(43.7-48.2)	(22-28)	(46.6-53.2)	(17-37)
14,774	27	13,910	21	55.7	24	56.1	35	46.9	23	47.8	34
(12,072-17,557)	(13-43)	(11,549-16,710)	(8-37)	(54.7-56.8)	(20-26)	(52.4-59.5)	(25-42)	(45.1-48.6)	(20-26)	(44.2-51.0)	(24-42)
15,361	41	15,099	41	53.0	27	55.2	40	44.3	27	46.4	40
(12,601-18,581)	(41)	(12,414-18,230)	(23-46)	(51.4-54.4)	(29-33)	(52.6-57.6)	(31-42)	(42.2-46.3)	(26-33)	(43.8-48.9)	(32-42)
15,508	43	14,614	37	52.4	29	60.6	20	44.0	28	51.2	21
(12,589-18,732)	(19-46)	(11,947-17,617)	(14-45)	(51.1-53.6)	(27-34)	(57.4-63.8)	(13-32)	(42.2-45.9)	(27-34)	(48.0-54.0)	(14-32)
13,713	10	12,723	3	46.7	45	60.9	19	40.2	45	52.5	16
(11,166-16,563)	(5-21)	(10,356-15,313)	(3-12)	(45.5-47.8)	(43-46)	(59.5-62.2)	(15-26)	(38.7-41.7)	(40-45)	(50.4-54.3)	(13-22)
14,366	18	14,462	33	60.0	15	59.8	28	50.8	14	49.9	28
(11,679-17,423)	(6-43)	(11,862-17,640)	(8-45)	(58.6-61.2)	(12-30)	(54.4-62.7)	(14-38)	(48.7-52.8)	(11-17)	(46.1-53.6)	(15-38)
14,365	17	13,291	9	57.4	20	62.5	15	48.5	20	53.3	15
(11,770-17,302)	(7-43)	(10,712-16,349)	(3-36)	(52.1-61.5)	(12-30)	(56.2-68.4)	(6-35)	(44.3-52.1)	(12-28)	(48.2-58.8)	(5-32)
14,555	21	13,298	7	60.6	13	64.9	8	50.9	13	55.3	9
(12,022-17,541)	(11-39)	(10,950-15,894)	(4-24)	(59.2-61.9)	(11-16)	(63.1-66.7)	(6-14)	(48.8-53.0)	(11-17)	(52.8-57.6)	(6-14)
14,811	28	14,194	28	52.1	31	59.4	25	43.7	31	50.2	25
(11,975-17,817)	(10-44)	(11,615-17,119)	(10-42)	(51.3-52.8)	(28-34)	(55.4-62.7)	(14-37)	(41.9-45.4)	(27-35)	(46.6-53.5)	(15-37)
14,670	26	14,148	25	50.9	33	56.9	32	42.9	33	48.2	32
(11,995-17,800)	(9-43)	(11,567-17,461)	(8-43)	(46.3-50.0)	(26-45)	(50.6-63.0)	(15-43)	(39.2-46.3)	(26-45)	(43.2-53.2)	(16-43)
13,184	5	12,884	4	63.0	7	64.8	9	53.8	6	55.5	8
(10,866-15,755)	(3-14)	(10,647-15,396)	(3-16)	(62.2-63.7)	(6-10)	(62.8-66.6)	(6-15)	(52.0-55.5)	(6-9)	(53.2-57.8)	(6-14)
13,503	7	15,483	44	60.9	12	47.5	45	52.0	11	40.3	45
(11,095-16,398)	(3-26)	(12,896-18,367)	(31-46)	(59.7-62.1)	(11-15)	(44.9-50.1)	(44-46)	(50.0-53.9)	(8-14)	(37.7-42.6)	(44-46)
15,298	38	15,200	42	49.6	38	57.2	33	41.4	39	47.7	36
(12,537-18,572)	(17-46)	(12,479-18,353)	(27-46)	(47.6-51.1)	(33-43)	(55.1-59.1)	(26-38)	(39.3-43.4)	(34-45)	(45.2-50.0)	(29-39)
14,116	14	14,145	26	56.1	22	63.6	12	47.6	22	53.8	13
(11,612-16,973)	(6-39)	(11,563-17,065)	(8-42)	(55.2-57.0)	(20-25)	(61.0-65.9)	(7-19)	(45.7-49.2)	(19-24)	(51.0-56.5)	(8-20)
15,082	35	14,549	36	48.7	42	52.9	42	41.0	42	45.0	41
(12,396-18,118)	(19-44)	(12,011-17,620)	(17-43)	(48.0-49.5)	(38-44)	(51.1-54.7)	(37-43)	(39.3-42.4)	(37-44)	(43.0-47.0)	(37-43)
14,609	24	13,991	24	48.7	43	57.3	31	41.1	41	48.6	31
(11,996-17,685)	(11-40)	(11,430-16,879)	(9-39)	(48.1-49.3)	(38-44)	(53.4-60.4)	(21-40)	(39.5-42.6)	(37-44)	(45.0-51.8)	(21-40)
15,248	36	14,307	29	60.2	14	64.5	11	50.3	15	54.3	11
(12,506-18,559)	(18-46)	(11,651-17,295)	(12-46)	(59.2-61.2)	(12-16)	(60.7-67.6)	(6-21)	(48.1-52.1)	(12-18)	(50.4-57.5)	(6-24)
11,333	1	10,953	1	68.9	2	73.2	2	60.3	1	63.9	1
(9,373-13,820)	(1-2)	(9,055-13,131)	(1-2)	(68.7-69.1)	(1-3)	(72.9-73.5)	(1-3)	(58.5-61.8)	(1-2)	(62.1-65.6)	(1-2)
15,342	39	14,841	39	50.0	36	52.6	43	41.8	38	44.5	43
(12,595-18,469)	(18-46)	(12,229-18,090)	(18-46)	(49.2-50.8)	(33-40)	(50.1-55.0)	(37-44)	(40.1-43.3)	(34-42)	(42.0-47.0)	(38-44)
13,774	11	13,809	14	62.4	9	61.6	17	53.2	8	52.5	17
(11,339-16,490)	(4-28)	(11,363-16,481)	(6-37)	(61.6-63.2)	(7-11)	(59.2-63.9)	(12-26)	(51.3-54.9)	(6-11)	(50.0-55.1)	(12-25)
14,526	19	13,915	22	46.2	46	57.8	30	38.9	46	44.9	30
(11,819-17,512)	(8-42)	(11,287-16,885)	(7-39)	(45.4-46.9)	(43-46)	(52.9-61.8)	(17-40)	(37.4-40.4)	(44-46)	(48.8-52.9)	(18-41)
14,273	15	13,842	16	55.1	25	59.6	24	46.4	25	50.5	24
(11,679-17,286)	(10-30)	(11,322-16,609)	(10-32)	(53.9-56.2)	(22-27)	(57.8-61.3)	(18-31)	(44.4-48.3)	(22-27)	(48.2-52.5)	(19-31)
15,064	32	13,895	20	49.7	37	64.6	10	42.2	35	54.7	10
(12,345-18,050)	(15-45)	(11,380-16,696)	(7-40)	(48.4-50.9)	(33-42)	(63.2-65.9)	(7-14)	(40.4-43.9)	(32-40)	(52.7-56.7)	(8-15)
13,532	9	13,252	8	63.5	6	70.2	4	53.8	7	59.6	3
(11,028-16,399)	(3-28)	(10,889-15,942)	(3-33)	(62.0-64.6)	(5-10)	(67.6-72.7)	(3-5)	(51.5-56.0)	(5-10)	(56.7-62.5)	(2-5)
13,914	13	13,379	10	58.8	16	65.3	7	49.9	16	55.7	7
(11,500-16,946)	(5-30)	(10,885-16,061)	(4-32)	(57.9-59.7)	(15-19)	(63.7-67.0)	(6-13)	(48.0-51.7)	(14-19)	(53.3-57.8)	(6-13)
11,681	2	11,259	2	66.7	4	66.1	6	58.3	3	58.1	5
(9,463-14,149)	(1-4)	(9,160-13,151)	(1-3)	(66.1-67.3)	(3-4)	(65.3-66.9)	(6-9)	(56.5-60.0)	(2-4)	(56.2-59.7)	(3-6)
15,322	40	15,221	43	48.9	40	58.7	29	40.9	43	49.1	29
(12,518-18,441)	(20-46)	(12,420-18,389)	(26-46)	(47.6-50.1)	(35-44)	(56.5-60.6)	(20-35)	(39.1-42.8)	(36-45)	(46.5-51.5)	(23-36)
14,350	16	14,546	35	51.6	32	56.2	34	43.9	32	47.6	33
(11,688-17,404)	(6-43)	(11,973-17,647)	(13-44)	(45.1-56.8)	(21-46)	(48.1-62.2)	(15-44)	(38.5-48.3)	(21-45)	(41.4-52.4)	(18-44)
12,905	3	13,826	15	64.6	5	59.9	23	55.5	5	51.1	22
(10,595-15,408)	(2-11)	(11,406-16,608)	(8-34)	(63.5-65.8)	(4-6)	(57.5-61.8)	(16-31)	(53.5-57.3)	(4-6)	(48.8-53.2)	(16-30)
15,884	44	14,781	38	62.1	11	68.8	5	51.4	12	57.1	6
(12,830-18,747)	(25-46)	(12,215-17,870)	(21-45)	(61.2-63.1)	(7-12)	(67.0-70.6)	(3-6)	(49.0-53.4)	(10-16)	(54.4-59.5)	(4-9)
13,819	12	15,781	46	62.3	10	49.5	44	53.0	9	41.9	44
(11,344-16,614)	(5-29)	(12,925-18,907)	(36-46)	(60.9-63.0)	(6-12)	(46.7-52.5)	(42-46)	(51.0-55.0)	(6-11)	(39.3-44.5)	(42-46)
14,553	20	14,177	27	56.7	21	61.7	16	47.7	21	52.2	18
(11,922-17,667)	(9-42)	(11,555-17,240)	(9-42)	(55.8-57.4)	(19-23)	(59.9-63.8)	(12-24)	(45.8-49.5)	(19-24)	(49.7-54.5)	(13-25)
14,577	22	13,973	23	58.3	17	60.2	22	49.0	19	51.0	23
(11,954-17,755)	(10-42)	(11,447-16,887)	(6-40)	(57.3-59.2)	(16-20)	(57.2-62.9)	(14-32)	(46.9-50.9)	(16-21)	(47.9-53.7)	(14-32)
15,596	45	13,882	18	50.4	35	60.4	21	42.2	36	51.4	20
(12,780-18,763)	(26-46)	(11,434-16,785)	(7-39)	(49.0-51.9)	(32-40)	(58.2-62.5)	(15-30)	(40.4-44.0)	(32-41)	(49.0-53.8)	(15-28)
15,030	31	13,732	13	50.6	34	55.9	37	42.7	34	47.8	35
(12,440-17,875)	(14-46)	(11,383-16,403)	(6-36)	(49.5-51.7)	(32-38)	(53.3-58.2)	(29-41)	(41.0-44.3)	(31-38)	(45.1-50.3)	(28-40)
13,058	33	15,659	45	63.0	8	53.2	41	52.7	10	44.8	42
(12,485-17,828)	(18-44)	(12,987-18,558)	(36-46)	(61.8-64.1)	(6-11)	(50.0-56.3)	(34-44)	(50.6-54.6)	(7-12)	(41.9-47.5)	(36-44)


CHANGES IN LEADING CAUSES OF DALYS BETWEEN 1990 AND 2010 FOR COUNTRIES IN SUB-SAHARAN AFRICA

In the following figures, pointed arrows indicate causes that have increased by a greater amount than shown on the x-axis. For more country data, explore IHME's data visualization tools online: www.ihmeuw.org/GBDcountryviz.


Shifts in leading causes of DALYs in Angola, 1990-2010




Shifts in leading causes of DALYs in Benin, 1990-2010




Shifts in leading causes of DALYs in Botswana, 1990-2010




Shifts in leading causes of DALYs in Burkina Faso, 1990-2010




Shifts in leading causes of DALYs in Burundi, 1990-2010




Shifts in leading causes of DALYs in Cameroon, 1990-2010




Shifts in leading causes of DALYs in Cape Verde, 1990-2010




Shifts in leading causes of DALYs in the Central African Republic, 1990-2010




Shifts in leading causes of DALYs in Chad, 1990-2010




Shifts in leading causes of DALYs in Comoros, 1990-2010




Shifts in leading causes of DALYs in Congo, 1990-2010




Shifts in leading causes of DALYs in Côte d'Ivoire, 1990-2010



Shifts in leading causes of DALYs in Democratic Republic of the Congo, 1990-2010




Shifts in leading causes of DALYs in Eritrea, 1990-2010




Shifts in leading causes of DALYs in Ethiopia, 1990-2010




Shifts in leading causes of DALYs in Gabon, 1990-2010




Shifts in leading causes of DALYs in The Gambia, 1990-2010




Shifts in leading causes of DALYs in Ghana, 1990-2010




Shifts in leading causes of DALYs in Guinea, 1990-2010




Shifts in leading causes of DALYs in Guinea-Bissau, 1990-2010




Shifts in leading causes of DALYs in Kenya, 1990-2010




Shifts in leading causes of DALYs in Lesotho, 1990-2010




Shifts in leading causes of DALYs in Liberia, 1990-2010




Shifts in leading causes of DALYs in Madagascar, 1990-2010




Shifts in leading causes of DALYs in Malawi, 1990-2010




Shifts in leading causes of DALYs in Mali, 1990-2010




Shifts in leading causes of DALYs in Mauritania, 1990-2010




Shifts in leading causes of DALYs in Mauritius, 1990-2010




Shifts in leading causes of DALYs in Mozambique, 1990-2010




Shifts in leading causes of DALYs in Namibia, 1990-2010




Shifts in leading causes of DALYs in Niger, 1990-2010




Shifts in leading causes of DALYs in Nigeria, 1990-2010




Shifts in leading causes of DALYs in Rwanda, 1990-2010




Shifts in leading causes of DALYs in São Tomé and Príncipe, 1990-2010



Shifts in leading causes of DALYs in Senegal, 1990-2010




Shifts in leading causes of DALYs in Seychelles, 1990-2010




Shifts in leading causes of DALYs in Sierra Leone, 1990-2010




Shifts in leading causes of DALYs in Somalia, 1990-2010




Shifts in leading causes of DALYs in South Africa, 1990-2010




Shifts in leading causes of DALYs in Sudan, 1990-2010




Shifts in leading causes of DALYs in Swaziland, 1990-2010




Shifts in leading causes of DALYs in Tanzania, 1990-2010




Shifts in leading causes of DALYs in Togo, 1990-2010




Shifts in leading causes of DALYs in Uganda, 1990-2010



Shifts in leading causes of DALYs in Zambia, 1990-2010



Shifts in leading causes of DALYs in Zimbabwe, 1990-2010





THE WORLD BANK



GBD

